

Access this article online

Quick Response Code:

Website:  
jecajournal.comDOI:  
10.4103/jeca.jeca\_21\_18

# Hepatoprotective effect of methanol fruit pulp extract of *Musa paradisiaca* on carbon tetrachloride-induced liver toxicity in Wistar rats

Mustapha Tosin Issa, Abel Nosereme Agbon, Sadiya Ufeli Balogun<sup>1</sup>, Onesimus Mahdi<sup>2</sup>, Khadijat Abubakar Bobbo<sup>2</sup>, Felix Olu Ayegbusi<sup>3</sup>

## Abstract:

**CONTEXT:** *Musa paradisiaca* (Banana) fruit pulp has been used in folk medicine to treat various kinds of ailments, such as dysentery, diarrhea, bronchitis, ulcer, fevers, and hemorrhages in different parts of the globe, including Nigeria, Western Africa.

**AIM:** This study was designed to histologically and biochemically assess the protective effect of methanol fruit pulp extract of *M. paradisiaca* (MFMP) on carbon tetrachloride (CCl<sub>4</sub>)-induced hepatotoxicity in Wistar rats.

**MATERIALS AND METHODS:** Twenty-four Wistar rats were divided into six groups (I–VI;  $n = 4$ ). Group I (control) was administered distilled H<sub>2</sub>O (2 ml/kg), whereas Groups II, III, and IV were administered MFMP (500 mg/kg, 1000 mg/kg, and 1500 mg/kg, respectively) and Group V administered Silymarin (100 mg/kg), as the reference drug, for a period of 14 days. Hepatotoxicity was induced in rats by the administration of CCl<sub>4</sub> (1 ml, 1:1 solution: olive oil). On the 15<sup>th</sup> day, Groups II–VI were administered single dose of CCl<sub>4</sub>. All administrations were through the oral route. After 12 h of CCl<sub>4</sub> administration, rats were euthanized and liver organs harvested for routine (H and E) histological tissue processing and blood samples collected for biochemical analysis of serum liver enzymes (alanine transaminase, aspartate transaminase, and alkaline phosphatase).

**RESULTS:** MFMP-treatment revealed remarkable histoarchitectural preservation of the liver parenchyma against CCl<sub>4</sub>-induced liver damage and decreased ( $P < 0.05$ ) serum liver enzyme levels elevated by CCl<sub>4</sub>. Hepatoprotective activity was comparable with that of the reference drug, Silymarin.

**CONCLUSION:** Result suggests that MFMP possesses hepatoprotective potentials against chemically-induced acute hepatotoxicity in Wistar rats. Hepatoprotective potential of MFMP is possible as a result of its antioxidant properties.

## Keywords:

Carbon tetrachloride, hepatoprotective, *Musa paradisiaca*, Wistar rat

## Introduction

Man is exposed to various exogenous compounds, such as drugs and pollutants in their daily activities. The liver plays a critical role in the process of chemical alteration of most of these compounds (Chiang, 2014; Ullah *et al.*,

2016). This physiological activity of the liver results in the generation of highly reactive free radicals in the cell, which bind to membrane lipids causing lipid peroxidation and consequently damage the liver (Ali *et al.*, 2014; Ye *et al.*, 2018). CCl<sub>4</sub> is an established hepatotoxin, frequently used as a model of experimental hepatotoxicity (Cheng *et al.*, 2013).

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**How to cite this article:** Issa MT, Agbon AN, Balogun SU, Mahdi O, Bobbo KA, Ayegbusi FO. Hepatoprotective effect of methanol fruit pulp extract of *Musa paradisiaca* on carbon tetrachloride-induced liver toxicity in wistar rats. J Exp Clin Anat 2018;17(1):1-7.

Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Ahmadu Bello University, Zaria, <sup>1</sup>Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Kogi State University, Ayingba, <sup>2</sup>Department of Human Anatomy, College of Medical Sciences, Gombe State University, Gombe, <sup>3</sup>Department of Chemical Pathology, Faculty of Clinical Sciences, College of Health Sciences, Ahmadu Bello University Teaching Hospital, Shika, Nigeria

## Address for correspondence:

Dr. Abel Nosereme Agbon, Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Health Sciences, Ahmadu Bello University, Zaria, Nigeria. E-mail: bellulu09@gmail.com

Side effect manifestation is a major challenge with the administration of conventional or synthetic drugs for the treatment of liver-related diseases (Toori *et al.*, 2015; Zarezade *et al.*, 2018). Thus, it is imperative to seek for new medicines for liver disease, especially those of natural origin which are considered to be effective and safe alternative treatments, readily available and accessible (Yao *et al.*, 2016; Gyawali *et al.*, 2017). *Musa paradisiaca* (banana) is a tree-like herb and various parts of the plant used in folk medicine for a variety of ailments (Enechi *et al.*, 2014; Abbas *et al.*, 2016). This study histologically and biochemically assessed the hepatoprotective effect of methanol fruit pulp extract of *M. paradisiaca* (MFMP) on CCl<sub>4</sub>-induced liver toxicity in Wistar rats. This study is imperative to scientifically demonstrate the pharmacological property of *M. paradisiaca* *in vivo* as potential therapeutics for inflammatory liver-related diseases.

## Materials and Methods

### Plant material collection and identification

*M. paradisiaca* (banana) fruit were obtained from a local market in Zaria, Kaduna State, Nigeria. Plant material was authenticated in the Herbarium Unit of the Department of Botany, Faculty of Life Sciences, Ahmadu Bello University (ABU), Zaria, with the Voucher Specimen Number: 173.

### Extraction of plant materials

Preparation of MFMP was conducted at the Department of Pharmacognosy and Drug Development, Faculty of Pharmaceutical Sciences, ABU, Zaria, employing the method of maceration. A brief description of the extraction protocol was as follows: shade-dried *M. paradisiaca* fruit pulp was pulverized, of which 250 g was macerated in 2.5 L of 100% methanol for 24 h. After which the solution was filtered using a Whatmann filter paper and the filtrate evaporated to dryness using H-H Digital Thermometer Water Bath (Mc Donald Scientific International-22050 Hz1.0A International Number) at 65°C. A yield of 60% of the extract was obtained.

### Experimental animals

Twenty-four adult Wistar rats (male and female, 170–200 g) were obtained from Human Anatomy Animal House, Faculty of Basic Medical, College of Health Sciences, ABU, Zaria, housed in new wired cages in the same facility and allowed to acclimatized for 2 weeks before the commencement of experiments. The rats were housed under standard laboratory condition, light and dark cycles of 12 h, and were provided with standard rodent pellet diet and water *ad libitum*. The rats were categorized into control and treatment groups.

### Drugs

- Olive oil (finest cold drawn) Bell, Sons and Co (DRUGGISTS) Ltd, Southport PR9 9AL, England
- Carbon tetrachloride (CCl<sub>4</sub>) May and Baker limited Dagenham, England
- Silymarin (Micro Labs Limited 92, Sipcot, Hosur-635 126 India).

### Experimental design

Twenty-four rats were divided into six Groups (I–VI) of four rats each. Group I was control, administered distilled H<sub>2</sub>O (2 ml/kg) while Groups II–VI were treatment groups. Liver toxicity was induced in rats by the administration of CCl<sub>4</sub> (1 ml, 1:1 solution: olive oil) as reported by Ikyembe *et al.* (2014). Groups II, III, and IV were administered MFMP (500 mg/kg, 1000 mg/kg and 1500 mg/kg, respectively) and Group V was administered Silymarin (100 mg/kg) as the reference drug, for a period of 14 days. After 12 h of the last MFMP or Silymarin administration; on the 15<sup>th</sup> day, all rats in the treatment groups (Groups II–VI) were administered a single dose of CCl<sub>4</sub>. All administrations were via the oral route. After 12 h of CCl<sub>4</sub> administration, rats were humanely sacrificed under chloroform anesthesia and liver harvested for routine (hematoxylin and Eosin, H and E, staining) histological tissue processing for light microscopy and blood samples collected in 5 ml plain sample bottles from jugular vein for biochemical analysis of liver serum enzymes (aspartate aminotransferase, aspartate transaminase [AST]; alanine aminotransferase, alanine transaminase [ALT], and alkaline phosphatase [ALP]) analysis.

### Phytochemical screening

Phytochemical analysis of MFMP was conducted in the Department of Pharmacognosy and Drug Development, Faculty of Pharmaceutical Sciences, ABU, Zaria. The method of Trease and Evans (2002) for phytochemical screening was adopted.

### Histological studies

The harvested liver organs of the rats were fixed in 10% buffered formalin. Tissues were processed routinely (H and E) for light microscopic examination in the Histology laboratory Unit of the Department of Human Anatomy, ABU, Zaria.

### Biochemical studies

Biochemical analysis of serum liver enzymes (AST, ALT, and ALP) were assayed in the Department of Chemical Pathology, ABU Teaching Hospital, Shika using assay kits according to the manufacturer's instructions.

### Data analysis

Results obtained were analyzed using the statistical software, Statistical Package for Social Scientist (SPSS version 18.0, SPSS Inc., 233 South Wacker Drive, 11

th Floor, Chicago, IL 60606-641, USA) and Microsoft Office Excel 2010 for charts. Results were expressed as mean  $\pm$  standard error of the mean and the presence of significant differences among means of the groups was determined using one-way ANOVA with least significance difference *post hoc test* for significance. Values were considered significant when value of  $P \leq 0.05$ .

## Results

### Phytochemical screening

Qualitative phytochemical analysis of MFMP produced positive reaction for each of the following secondary metabolites: carboxylic acid, tannins, saponin, flavonoids, terpenoids, and steroid, whereas anthroquinone and glycosides were absent.


### Histological studies

Light microscopic examination of liver sections of the rats in the control group revealed normal histoarchitecture of the liver parenchyma; the characteristic appearance of hepatic lobule unit: centrilobular venules (central veins), array of interconnected plates of hepatocytes (constituting two-thirds of the mass of the liver) radiating from the central vein, separated by vascular spaces (sinusoids) and portal tract/triad (hepatic portal vein, hepatic vein, and bile duct) [Figure 1a].

The liver sections of the rats treated with  $\text{CCl}_4$  only (Group VI) revealed severe histoarchitectural distortion of the liver parenchyma which manifested as diffused necrosis of hepatocytes with pyknotic nuclei and hepatocellular vacuolation, infiltration of inflammatory cells and central vein congestion [Figure 1b]. The liver sections of Silymarin +  $\text{CCl}_4$ -treated rats showed normal cytoarchitecture of the liver with mild histoarchitectural distortion; infiltration of inflammatory cells and localized hepatocellular vacuolations, when compared with the severe distortions observed with  $\text{CCl}_4$  only-treated group [Figure 1c]. The liver sections of MFMP (500 mg/kg) +  $\text{CCl}_4$  and MFMP (1000 mg/kg) +  $\text{CCl}_4$ -treated rats, showed normal cytoarchitecture with mild histoarchitectural distortion, such as sinusoidal dilatation and infiltration of inflammatory cells [Figure 1d and e]. In MFMP (1500 mg/kg) +  $\text{CCl}_4$ -treated rats, the liver sections of rats revealed histoarchitectural distortion of liver parenchyma manifesting as hepatocellular necrosis, vacuolations, and sinusoidal dilations. However, observed distortions were not as severe when compared to  $\text{CCl}_4$ -treated group [Figure 1f].


### Biochemical studies

Serum liver enzymes (AST, ALT, and ALP) were analyzed and revealed the following: relative to the control, elevated levels of AST were observed in all the treated groups, especially ( $P < 0.05$ ) in  $\text{CCl}_4$ -treated group [Figure 2]. No remarkable ( $P > 0.05$ ) difference




**Figure 1:** Micrograph of liver sections of Wistar rat. H and E, (Mag  $\times 100$ ). (a) Control (2 ml/kg distilled  $\text{H}_2\text{O}$ ), showing normal histology. Central vein (c); Hepatocyte (h); Sinusoid (s). (b)  $\text{CCl}_4$ -treated, showing histoarchitectural distortion. Congested central vein (cC); Necrosis (n); Inflammatory cell infiltration (i); Hepatocellular vacuolation (v). (c) Silymarin (100 mg/kg) +  $\text{CCl}_4$ -treated, showing mild distortion of the histoarchitecture of the liver. Inflammatory cells infiltration (i); Necrosis (n). (d), Methanol fruit pulp extract of *Musa paradisiaca* (500 mg/kg) +  $\text{CCl}_4$ -treated, showing mild histoarchitectural distortion of the liver. Central vein (c); Inflammatory cells infiltration (i); Sinusoid (s). (e): Methanol fruit pulp extract of *Musa paradisiaca* (1000 mg/kg) +  $\text{CCl}_4$ -treated, showing mild distortion of the histoarchitecture of the liver. Central vein (c); Sinusoids (s); Inflammatory cells infiltration (i). (f) Methanol fruit pulp extract of *Musa paradisiaca* (1500 mg/kg) +  $\text{CCl}_4$ -treated, showing mild distortion of the histoarchitecture of the liver. Central vein (c); Necrosis (n); Sinusoidal dilation (s); Hepatocellular vacuolation (v)





**Figure 2:** Effect of methanol fruit pulp extract of *Musa paradisiaca* on serum aspartate transaminase levels in Wistar rats.  $n = 4$ ; means  $\pm$  standard error of the mean. One-way ANOVA least significant difference *post hoc* test: \* =  $P < 0.05$  significant difference when compared with the control. Control: 2 ml/kg (distilled water); MPLo, MPMi and MPHi (methanol fruit pulp extract of *Musa paradisiaca* 500 mg/kg, 1000 mg/kg and 1500 mg/kg, respectively); SIL: Silymarin (100 mg/kg); CCl<sub>4</sub>: Carbon tetrachloride (1 ml)



**Figure 3:** Effect of methanol fruit pulp extract of *Musa paradisiaca* on serum alanine transaminase levels in Wistar rats.  $n = 4$ ; means  $\pm$  standard error of the mean. One-way ANOVA least significant difference *post hoc* test:  $P > 0.05$  when compared with the control. Control: 2 ml/kg (distilled water); MPLo, MPMi, and MPHi (methanol fruit pulp extract of *Musa paradisiaca* 500 mg/kg, 1000 mg/kg, and 1500 mg/kg, respectively); SIL: Silymarin (100 mg/kg); CCl<sub>4</sub>: Carbon tetrachloride (1 ml)

in levels of ALT was observed in all the treated groups when compared to the control [Figure 3].

Remarkable ( $P < 0.01$ ) up-regulation of ALP level was observed in all the treated groups, except MFMP (1,500 mg/kg) + CCl<sub>4</sub> and Silymarin (100 mg/kg) + CCl<sub>4</sub>-treated groups when compared to the control. ALP levels of MFMP (1,500 mg/kg) + CCl<sub>4</sub> and Silymarin (100 mg/kg) + CCl<sub>4</sub>-treated groups were remarkably ( $P < 0.05$ ) decreased relative to CCl<sub>4</sub> only treated group [Figure 4].

## Discussion


In this study, preliminary qualitative phytochemical screening of MFMP was carried out and hepatoprotective activity was evaluated employing histological examinations of liver sections and biochemical analysis of serum liver enzymes of CCl<sub>4</sub>-induced hepatotoxicity model in Wistar rats.

In this study, phytochemical analysis of the plant extract, MFMP, revealed the presence of secondary metabolites such as, flavonoids, tannins terpenoids and steroid, and saponin, which is in accordance to reported constituents present in various parts of the plant (Abbas *et al.*, 2016; Sundaram *et al.*, 2018). Phytochemical constituents of natural agents with antioxidant properties have been reported to play pertinent roles in hepatoprotection from noxious exogenous substances (Swathi *et al.*, 2011; Sarian *et al.*, 2017). The presence of phenolic compound, such as flavonoid, has been established to have free radical scavenging and antioxidant activities (Heim *et al.*, 2002; Seyoum *et al.*, 2006).

CCl<sub>4</sub> is an established potent hepatotoxic compound, frequently used as a model of experimental hepatotoxicity

(Ahsan *et al.*, 2009; Cheng *et al.*, 2013) which presents with features analogous to human acute hepatitis (Li *et al.*, 2015) and a useful model system for assessing of hepatoprotective activity of plant agents (Adewusi and Afolayan, 2010; Huang *et al.*, 2018). Several researches have demonstrated the critical role of oxidative stress in the pathophysiology of CCl<sub>4</sub>-induced hepatotoxicity (Kepekci *et al.*, 2013; Hafez *et al.*, 2014). Within biological system, CCl<sub>4</sub> undergoes enzymatic activation, majorly CYP2E<sub>1</sub>, into the trichloromethyl free radical (CCl<sub>3</sub>) within the endoplasmic reticulum membrane, which is followed by chloromethylation, saturation, peroxidation, and progressive destruction of the unsaturated fatty acid of the endoplasmic reticulum membrane phospholipids (Peng *et al.*, 2009; Abbas *et al.*, 2016). These processes are known as lipid peroxidation, leading to functional, and structural disruption of hepatocytes (Hogade *et al.*, 2010; Ali *et al.*, 2014).

In this study, liver parenchymal histoarchitectural distortion consequent to CCl<sub>4</sub> administration manifested as severe hepatocellular necrosis and vacuolation, infiltration of inflammatory cells and central vein congestion. These are an indication of oxidative damage (Padhy *et al.*, 2007). Findings are in accordance with the reports of previous researches on histopathological changes following CCl<sub>4</sub> treatment in rodents (Mahli *et al.*, 2015; Huda and Mosaddik, 2018). Necrosis is a pathological form of cell death sequential to exposure to abnormal stressors, such as chemical injury or toxin. Necrotic cells, unable to maintain membrane integrity, leak out their content which could elicit inflammation in the surrounding tissue (Kumar *et al.*, 2009). Inflammation, infiltration of the inflammatory cells relative to CCl<sub>4</sub> intoxication as observed, is a protective response targeted to rid the biological system of cell



**Figure 4:** Effect of methanol fruit pulp extract of *Musa paradisiaca* on serum alkaline phosphatase levels in Wistar rats.  $n = 4$ ; means  $\pm$  standard error of the mean. One way ANOVA least significant difference *post hoc* test: \* =  $P < 0.05$  difference when compared with the control. a =  $P < 0.05$  difference when compared with  $CCl_4$ . Control: 2ml/kg (distilled water); MPLo, MPMi and MPHi (methanol fruit pulp extract of *Musa paradisiaca* 500 mg/kg, 1,000 mg/kg and 1,500 mg/kg, respectively); SIL: Silymarin (100 mg/kg);  $CCl_4$ : Carbon tetrachloride (1 ml)

injury inducing factors (e.g., toxin) and consequence of such injury (necrotic cells and tissues) (Kumar *et al.*, 2009). Congestion of central vein relative to  $CCl_4$  intoxication as observed, is a local increase in the volume of blood as a consequence of impaired outflow from the hepatic tissue (Cotran *et al.*, 2005).

Silymarin is an established antioxidant used in the evaluation of hepatoprotective activity of potential nutraceuticals (Flora *et al.*, 1998; Shalan *et al.*, 2005; Pradhan and Girish, 2006; Shaker *et al.*, 2010). Mild histoarchitectural changes in the liver sections observed in Silymarin-treated rats is indicative of the ameliorative effect of Silymarin against chemically-induced hepatotoxicity. Silymarin has anti-inflammatory potentials, act as a radical scavenger, thus, protecting membrane permeability and alter drug-induced histopathological changes (Song *et al.*, 2006; Mahli *et al.*, 2015).

The development and progression of liver-related diseases has been described in relation to free radicals generation and oxidative stress (Zhu *et al.*, 2012; Jadeja *et al.*, 2017). Thus, oxidative liver injury is commonly treated with the use of antioxidants (Sundaram *et al.*, 2018). Mild histoarchitectural changes; histo-parenchymal preservation of the liver sections observed in MFMP-treated rats, especially with 1000 mg/kg dose, is suggestive of the protective effect of MFMP against chemically-induced hepatotoxicity. Consumption of antioxidants is an essential means of preventing or delaying the appearance of liver diseases (Ganesan *et al.*, 2018). Mahmood *et al.* (2010) and Nirmala *et al.* (2012) reported the presence of antioxidants, tannins, and flavonoids, in *M. paradisiaca* to provide plant's hepatoprotective activity. Flavonoid

have been reported as antioxidant in plants, having free radicals scavenging activity and is able to reduce the formation of free radical, chelate metal catalysts, down-regulate alpha-tocopherol radicals, inhibit oxidases, and up-regulate the levels of antioxidant enzymes in plasma (Pietta, 2000; Heim *et al.*, 2002).

Plasma concentration levels of liver enzymes are important biochemical indicators of liver functionality (Zhao *et al.*, 2018). During hepatic damage, cellular enzymes such as AST, ALT, and ALP and bilirubin leak into the serum resulting in up-regulation of their serum concentration.  $CCl_4$  is known to induce hepatic alteration as, marked up-regulation in serum levels of ALP, AST and especially ALT which is considered the primary and specific marker of liver injury (Vozarova *et al.*, 2002; Anand *et al.*, 2011). In this study, remarkable up-regulation in the serum levels of AST and ALP indicated treatment-related hepatocellular damage in  $CCl_4$  (only)-treated rat. Findings are consistent with reports on elevated hepatic biomarker levels following  $CCl_4$  administration (Mosa and Khalil, 2015; Chen *et al.*, 2018). Critical to the efficacy of any hepatoprotective agent is, its capacity to either ameliorate the harmful effect or restore the normal hepatic physiology altered by a hepatotoxin (Palanivel *et al.*, 2008). Remarkable down-regulation of hepatic biomarker levels, especially ALP levels, in Silymarin-and MFMP-treated groups when compared to that of  $CCl_4$  (only)-treated rat is suggestive of treatment-related hepatoprotective activity. Zarezade *et al.* (2018) reported oral administration of Silymarin (100 mg/kg) showed a significant decrease in the serum levels of hepatic biomarkers. Findings are in accordance with hepatoprotective activities of natural agents employed in folk medicine. Several reports on the potential benefits of medicinal plants have demonstrated the protective effects of these plants' extracts on experimentally-induced hepatotoxicity (Osman *et al.*, 2011; Gong *et al.*, 2012; Ikyembe *et al.*, 2014). Mosa and Khalil (2015) reported a reduction of hepatic biomarkers levels in rats fed 10% peels of *M. paradisiaca*-supplemented diet. MFMP hepatoprotective effect could be mediated by its constituent phytochemical antioxidant properties targeted toward plasma membrane stabilization, thus preserving the structural integrity of hepatocytes; influence regeneration of damaged hepatocytes and repair of  $CCl_4$ -induced hepatic tissue damage (Osman *et al.*, 2011). Natural agents' ability to attenuate toxin-induced hepatotoxicity is related to their intrinsic antioxidant properties, such as flavonoids and saponins (Yoshikawa *et al.*, 2003; Prakash *et al.*, 2008) which could have played a critical role in the hepatoprotective activity of MFMP.

## Conclusion

MFMP possess potential hepatoprotective activity against  $CCl_4$ -induced hepatotoxicity that manifested as

histo-parenchymal preservation of liver sections and decrease in serum liver enzymes levels in Wistar rats. Hepatoprotective activity is probably consequent to the antioxidant activities of constituent phytochemicals.

### Financial support and sponsorship

Nil.

### Conflicts of interest

There are no conflicts of interest.

## References

1. Abbas K., Rizwani G.H., Zahid H., Javed T.M. (2016). Evaluation of hepatic activity of various morphological parts of *Musa paradisiaca* L. Afr J Pharm Pharmacol 10 (19):419-29.
2. Adewusi E.A., Afolayan A.J. (2010). A review of natural products with hepatoprotective activity. J Med Plants Res 4:1318-34.
3. Ahsan M.R., Islam K.M., Bulbul I.J., Musaddik M.A., Haque E. (2009). Hepatoprotective activity of methanol extract of some medicinal plants against carbon tetrachloride-induced hepatotoxicity in rats. Euro J Sci Res 37:302-10.
4. Ali H., Kabir N., Muhammad A., Shah M.R., Musharraf S.G., Iqbal N., Nadeem S. (2014). Hautriwaic acid as one of the hepatoprotective constituent of *Dodonaea viscosa*. Phytomed 21:131-40.
5. Anand K.V., Anandhi R., Pakkiyaraj M., Geraldine P. (2011). Protective effect of Chryson on carbon tetrachloride (CCl<sub>4</sub>)-induced tissue injury in male wistar rats. Toxicol Ind Health 27 (10):923-33.
6. Chen Y.J., Chou P., Hsu C.L., Hung J., Wu Y., Lin J. (2018). Fermented citrus lemon reduces liver injury induced by carbon tetrachloride in rats. Evid Based Complement Alternat Med 2018:1-10.
7. Cheng N., Ren N., Gao H., Lei X., Zheng J., Cao W. (2013). Antioxidant and hepatoprotective effects of *Schisandra chinensis* pollen extract on CCl<sub>4</sub>-induced acute liver damage in mice. Food Chem Toxicol 55:234-40.
8. Chiang J. (2014). Liver physiology: Metabolism and detoxification. In: McManus L.M., Mitchell R.N., editors. Pathobiology of Human Disease. Elsevier, San Diego, 1770-82.
9. Cotran R.S., Kumar V., Fausto N., Robbins S.L., Abbas A.K. (2005). Robbins and Cotran Pathologic Basis of Disease. 7<sup>th</sup> ed. Elsevier Saunders, St. Louis, MO, 878.
10. Enechi O.C., Odo C.E., Agosi P.O. (2014). Antioxidant vitamins, phytochemicals and proximate composition of the ethanol extract of the leaves of *Musa paradisiaca*. Afr J Pharm Pharmacol 8 (18):464-8.
11. Flora K., Hahn M., Rosen H., Benner K. (1998). Milk thistle (*Silybum marianum*) for the therapy of liver disease. Am J Gastroenterol 93:139-43.
12. Ganesan K., Jayachandran M., Xu B. (2018). A critical review on hepatoprotective effects of bioactive food components. Crit Rev Food Sci Nutr 58 (7):1165-229.
13. Gong F., Yin Z., Xu Q., Kang W. (2012). Hepatoprotective effect of *Mitragyna rotundifolia* Kuntze on CCl<sub>4</sub>-induced acute liver injury in mice. Afr J Pharm Pharmacol 6:330-5.
14. Gyawali R., Shrestha A., Khanal A., Pyakurel J.S., Joshi N., Bajaj P., Chaudhary P., Thapa R. (2017). Hepatoprotective properties of selected plants against paracetamol-induced hepatotoxicity in mice. J Instit Sci Technol 22 (1):147-51.
15. Hafez M.M., Al-Shabanah O.A., Al-Harbi N.O., Al-Harbi M.M., Al-Rejaie S.S., Alsurayea S.M., Sayed-Ahmed M.M. (2014). Association between paraoxonases gene expression and oxidative stress in hepatotoxicity induced by CCl<sub>4</sub>. Oxid Med Cell Longev 2014:1-12.
16. Heim K.E., Tagliaferro A.R., Bobilya D.J. (2002). Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. J Nutr Biochem 13:572-84.
17. Heim K.E., Tagliaferro A.R., Bobilya D.J. (2002). Flavonoid antioxidants; chemistry, metabolism and structure-activity relationships. J Nutr Biochem 13:572-84.
18. Hogade M.G., Patil K.S., Wadkar G.H., Mathapati S.S., Dhumal P.B. (2010). Hepatoprotective activity of *Morus alba* (Linn.) leaves extract against carbon tetrachloride induced hepatotoxicity in rats. Afr J Pharm Pharmacol 4:731-4.
19. Huang Z., Chen P., Wei-Wei S., Yong-Gang W., Hao W., Wei P., Li P.B. (2018). Antioxidant activity and hepatoprotective potential of quercetin 7-rhamnoside *in vitro* and *in vivo*. Molecules 23 (1188):1-13.
20. Huda M.N., Mosaddik A. (2018). Evaluation of hepatoprotective effect of a polyherbal formulation against carbon tetrachloride-induced hepatotoxicity in rats. Altern Integr Med 7 (2):1-4.
21. Ikyembe D., Pwavodi C., Agbon A.N. (2014). Hepatoprotective effect of methanolic leaf extract of *Anacardium occidentale* (cashew) on carbon-tetrachloride-induced liver toxicity in wistar rats. Sub Saharan Afr J Med 1:124-31.
22. Jadeja R.N., Devkar R.V., Nammi S. (2017). Oxidative stress in liver diseases; pathogenesis, prevention, and therapeutics. Oxid Med Cell Longev 2017:1-12.
23. Kepekci R.A., Polat S., Celik A., Bayat N., Saygideger S.D. (2013). Protective effect of *Spirulina platensis* enriched in phenolic compounds against hepatotoxicity induced by CCl<sub>4</sub>. Food Chem 141:1972-9.
24. Kumar V., Alla S.R., Krishnan K.S., Ramaswami M. (2009). Syndapin is dispensable for synaptic vesicle endocytosis at the *Drosophila larval* neuromuscular junction. Mol Cell Neurosci 40 (2):234-41.
25. Li C., Yi L.T., Geng D., Han Y.Y., Weng L.J. (2015). Hepatoprotective effect of ethanol extract from *Berberis lineate* against CCl<sub>4</sub>-induced acute hepatotoxicity in mice. Pharm Biol 53:767-72.
26. Mahli A., Koch A., Czech B., Peterburs P., Lechner A., Haunschild J., Muller M., Hellerbrand C. (2015). Hepatoprotective effect of oral application of a silymarin extract in carbon tetrachloride induced hepatotoxicity in rats. Clin Phytosci 1:5.
27. Mahmood A., Omar M.N., Ngah N. (2010). Phytochemical constituents and antioxidant activities of banana flower of *Musa paradisiaca*. In: 3<sup>rd</sup> International Conference on Advancement in Science and Technology (iCAST) 27-29 November 2010, Vistana Hotel, Kuantan, Pahang.
28. Mosa Z.M., Khalil A.F. (2015). The effect of banana peels supplemented diet on acute liver failure rats. Ann Agr Sci 60 (2):373-9.
29. Nirmala M., Girija K., Lakshman K., Divya T. (2012). Hepatoprotective activity of *Musa paradisiaca* on experimental animal models. Asian Pac J Trop Biomed 2:11-5.
30. Osman M., Ahmed M., Mahfouz S., Elaby S. (2011). Biochemical studies on the hepatoprotective effects of pomegranate and guava ethanol extracts. NY Sci J 4:27-41.
31. Padhy B.M., Srivastava A., Kumar V. (2007). *Calotropis procera latex* affords protection against carbon tetrachloride -induced hepatotoxicity in rats. J Ethnopharm 113 (3):498-502.
32. Palanivel M.G., Raj Kapoor B., Kumar R.S., Einstein J.W., Kumar E.P., Kumar M.R., Kavitha K., Kumar M.P., Jayakar B. (2008). Hepatoprotective and antioxidant effect of *Pisonia aculeata* L. against CCl<sub>4</sub> induced hepatic damage in rats. Sci Pharm 76:203215.
33. Peng C., Chunying L., Wenqing P., Yue Z., Wei D., Shiming W., Zhang J. (2009). The protective role of per 2 against carbon tetrachloride induced hepatotoxicity. Am J Pathol 174:63-70.
34. Pietta P.G. (2000). Flavonoids as antioxidants. J Nat Prod 63:1035-42.
35. Pradhan S.C., Girish C. (2006). Hepatoprotective herbal drug,

- silymarin from experimental pharmacology to clinical medicine. *Indian J Med Res* 124:491-504.
36. Prakash T., Fadadu S.D., Sharma U.R., Surendra V., Goli D., Stamina P., Kotresha D. (2008). Hepatoprotective activity of leaves of *Rhododendron arboreum* in CCl<sub>4</sub> induced hepatotoxicity in rats. *J Med Plants Res* 2:315-20.
  37. Sarian M.N., Ahmed Q.U., So'ad S.Z., Alhassan A.M., Murugesu S., Perumal V., Syed Mohamad S.N., Khatib A., Latip J. (2017). Antioxidant and antidiabetic effects of flavonoids: A structure-activity relationship based study. *Biomed Res Int* 2017:1-14.
  38. Seyoum A., Asres K., El-Fiky F.K. (2006). Structure-radical scavenging activity relationships of flavonoids. *Phytochem* 67:2058-70.
  39. Shaker E., Mahmoud H., Mnaa S. (2010). Silymarin, the antioxidant component and *Silybum marianum* extracts prevents liver damage. *Food Chem Toxicol* 48:803-6.
  40. Shalan M.G., Mostafa M.S., Hassouna M.M., El-Nabi S.H., El-Refai A. (2005). Amelioration of lead toxicity on rat liver with Vitamin C and silymarin supplements. *Toxicol* 206:1-15.
  41. Song Z., Deaciuc I., Song M., Lee D.Y., Liu Y., Ji X., McClain C. (2006). Silymarin protects against acute ethanol induced hepatotoxicity in mice. *Alcohol Clin Exp Res* 30:407-13.
  42. Sundaram I.K., Sarangi D.D., Sundararajan V., George S., Mohideen S.S. (2018). Poly herbal formulation with anti-elastase and anti-oxidant properties for skin anti-aging. *BMC Compl Altern Med* 18:33.
  43. Swathi D., Jyothi B., Sravanthi C. (2011). A review: Pharmacognostic studies and pharmacological actions of *Musa paradisiaca*. *Int J Innov Pharm Res* 2 (2):122-5.
  44. Toori M.A., Joodi B., Sadeghi H., Jafari M., Talebianpoor M.S., Mehraban F., Mostafazadeh M., Ghavamizadeh M. (2015). Hepatoprotective activity of aerial parts of *Otostegia persica* against carbon tetrachloride-induced liver damage in rats. *Avicenna J Phytomed* 5 (3):238-46.
  45. Trease G.E., Evans W.C. (2002) *Phytochemicals in: Pharmacognosy*. 15<sup>th</sup> ed. W.B. Sanders Publishers, London.
  46. Ullah I., Khan J.A., Adhikari A., Shahid M. (2016). Hepatoprotective effect of *Monothecha buxifolia* fruit against antitubercular drugs-induced hepatotoxicity in rats. *Bangladesh J Pharmacol* 11:248-56.
  47. Vozarova B., Stefan N., Lindsay R.S., Saremi A., Pratley R.E., Bogardus C., Tataranni P.A. (2002). High alanine aminotransferase is associated with decreased hepatic insulin sensitivity and predicts the development of type 2 diabetes. *Diabetes* 51:1889-95.
  48. Yao H., Qiao Y.J., Zhao Y.L., Tao X.F., Xu L.N., Yin L.H., Qi Y, Peng J.Y. (2016). Herbal medicines and nonalcoholic fatty liver disease. *World J Gastroenterol* 22 (30):6890-905.
  49. Ye H., Nelson L.J., Gómez del Moral M., Martínez-Naves E., Cubero F.J. Qi Y., Peng J.Y. (2018). Dissecting the molecular pathophysiology of drug-induced liver injury. *World J Gastroenterol* 24 (13):1373-85.
  50. Yoshikawa M., Morikawa T., Kashima Y., Ninomiya K., Matsuda H. (2003). Structures of new dammarane-type triterpene saponins from the flower buds of *Panax notoginseng* and hepatoprotective effects of principal Ginseng saponins. *J Nat Prod* 66:922-7.
  51. Zarezade V., Moludi J., Mostafazadeh M., Mohammadi M., Veisi A. (2018). Antioxidant and hepatoprotective effects of *Artemisia dracunculus* against CCl<sub>4</sub>-induced hepatotoxicity in rats. *Avicenna J Phytomed* 8 (1):51-62.
  52. Zhao Z., Chang J., Lin L., Tsan F., Chang H., Wu C. (2018). Comparison of the hepatoprotective effects of four endemic *Cirsium* species extracts from Taiwan on CCl<sub>4</sub>-induced acute liver damage in C57BL/6 mice. *Int J Mol Sci* 19 (1329):1-18.
  53. Zhu R.Z., Wang Y.J., Zhang L.Q., Guo Q.L. (2012). Oxidative stress and liver disease. *Hepatol Res* 42:741-9.